

CHANGE OF BIOACTIVE CONTENT IN SEED PROPAGATED TEA PLANTS AT DIFFERENT SHOOTING PERIODS

Neva Karatas ¹, Sezai Ercisli ², Liidia Samarina ³

¹ Ataturk University Faculty of Health Sciences
Department of Nutrition and Dietetics,
Erzurum, Turkey

² Ataturk University Faculty of Agriculture,
Department of Horticulture,
Erzurum, Turkey; e-mail: sercisli@gmail.com

³ Federal Research Centre
the Subtropical Scientific Centre of the Russian Academy of Sciences,
Biotechnology Department,
Sochi, Russia, e-mail: q11111w2006@ya.ru

Tea production is vital for east Black Sea region for farmers because it is a dominant agricultural product. The majority of tea gardens in the region were multiplied by farmers using seeds in the last 50 years resulted a great variation for most of the plant characteristics. The present study describes variation in shooting periods on the bioactive content of seed propagated tea plant. Results indicated significant differences among most of the searched parameters among shooting periods. Total phenolic content was the highest at the first shooting period (between 6.45–7.03 g gallic acid equivalent/100 g) dry weight (DW) basis. Among the catechin, Epigallocatechin gallate (EGCG) was found to be dominant in black tea samples (0.756–0.941 g/100 g DW) and followed by epigallocatechin (0.590–0.752 g/100 g DW) and epicatechine gallate (0.510–0.654 g/100 g DW), respectively. In general, first shooting period (May) gave the highest catechin content that improved tea quality and with following shooting periods catechin content decreased steadily. Total phenolic content was found between 6.45–7.03 g GAE/100 g DW among shooting periods. The results suggested that first shooting period of black tea samples had higher bioactive content.

Key words: tea, diversity, phytochemicals, seed propagation.

Tea is one of the most important beverages in the world and black tea production in the world increased steadily and reached 4.8 million tons in recent years. On an average, during the last two decades in the production front, China contributed 35 % of the world tea production and followed by India (21 %), Kenya (8 %) and Sri Lanka (7 %). Turkey is the 5th biggest tea producer in the world with 225.000 tons black tea production per year. The country shares 5 % world tea production [3].

Tea (*Camellia sinensis*) is the most consumed drink in the world except for water, owing to its pleasant sensory properties, broad health benefits and unique sociocultural characteristics. Usually, tea could be classified into six categories, i.e., green tea (unfermented), yellow tea (slight-fermented), white tea (mild-fermented), oolong tea (semi-fermented), black tea (fermented) and dark tea (post-fermented) according to the varying fermentation extent [17].

At the start of tea cultivation in Turkey, the primary goal was to meet the domestic demand only. Today, Turkey holds a significant place among the world's largest tea producers. But still great percentage of products is marketed in domestic market. All of the tea plantation areas are located in the eastern part of Black Sea Region. Tea cultivation in Turkey contributes significantly to the economic income of local farmers in the east Black Sea region [2].

The main tea production area in Turkey is Rize province with 50.583 hectares area and followed by Trabzon (15.328 hectares), Artvin (8 430 hectares) and Giresun-Ordu (2 299 hectares). Approximately 66 % of total tea plantation areas located in Rize province whilst the rest of them are located in Trabzon, Artvin, Giresun and Ordu respectively [13].

Tea growing areas of Turkey is far away from Ecuador line compared the other most important the producers such as China, Kenya, India, Sri Lanka etc. indicating great differences from the rest of the countries. Because temperature does not falls up to minus degree in tea production areas and tea production is covering the whole year in those countries. However in Turkey, tea plantations active 6 months and snowfalls on Turkish tea plantations in winter months bring them an extra important characteristic. Tea plants harvested in in general 3 times in May, July and September, respectively. Shooting (harvesting) period, which is specific to tea cultivation in Turkey, significantly affected almost all of the quality parameters of black tea. Due to winter dormancy and short cultivation period, Turkish tea plantations are pesticide free and Turkish tea called 'natural tea' compared to other tea produced in the world [6, 15].

Tea was introduced into Turkey with the efforts of Zihni Derin and Ali Rıza Erten at the end of the 1930s. The most of the tea plantations (90–95 %) established by seeds and this tradition still have ongoing. Continuous seed propagation in Turkey has produced populations with different yield and quality properties, reflecting wide genetic variation. Thus, clonal selection studies were previously conducted on those germplasms in the Black Sea region and several promising tea clones such as 'Tuglali-10', 'Derepazari-7', and 'Pazar-20' have been released [10]. Unfortunately these clones have not become widespread so far.

Previous studies clearly revealed that black tea had one of the richest sources of bioactive phytochemicals including theaflavins, thearubigins, catechins, and phenolics [8, 11, 12].

The aim of the current study was to evaluate and compare the bioactive content of seed propagated tea populations to reveal if there are any differences among populations.

Material and Methods. The fresh tea leaves (*Camellia sinensis* var. *sinensis*) were harvested as a group of two or three leaves and a bud or mature leaves without a bud during three consecutive shooting periods (May, July, and September) in 2014 from a single seed propagated tea population found in tea garden in Ikizdere town belongs to Rize province located on the Black Sea coast in Turkey. After plucking, the leaves were immediately processed into black tea according to the Turkish Tea Board standards.

Preparation of extract. Phenolic extracts were prepared according to Zuo et al. [18]. First 50 mL of aqueous methanol (80 %) containing 0.15 % HCl was poured onto 2 g of tea sample, and the phenolic compounds were extracted at 40 °C for 3 h and filtered through coarse filter paper. The filtrate was used for the total phenolic content determination by Folin–Ciocalteu assay and the filtrate was further filtered through 0.45 µm of nylon membrane filter before HPLC analysis [11].

Total phenolic content. The total phenolic content was determined by the Folin–Ciocalteu method [1]. The results were expressed as mg gallic acid equivalent (GAE) per 100 g dry weight (DW) basis.

Catechins content. The catechin group including epigallocatechin gallate (EGCG), epigallocatechin (EGC), epicatechin (EC), epicatechin gallate (ECG), catechin gallate (CG), gallic catechin gallate (GCG), catechin (C) and caffeine were determined by HPLC. The chromatographic separation, described by Zuo et al. [18], was used with some modifications in the mobile phase and UV detection. The absorbance of the catechins (EGCG, EGC, EC, ECG, CG, GCG and C) were determined at 280 nm and the results were expressed g/100 g dry weight (DW) basis.

Statistical Analysis. All data were analyzed using SPSS software and procedures. Analysis of variance tables were constructed using the Least Significant Difference (LSD) method at $p < 0.05$.

Results and Discussion. Table 1 shows individual catechin content of the black tea samples obtained from 3 shooting periods (May, July and September).

As indicated in Table 1, there were statistically significant differences among shooting periods on individual catechins except CG (tbl. 1). All of the catechins decreased significantly from the first to the third shooting periods. According to obtained results Epigallocatechin gallate (EGCG) was the predominant catechin in black tea samples and EGCG content was the highest at first shooting period (0.941 g/100 g DW), and followed by second shooting period (0.855 g/100 g DW) and finally it was the lowest at the last shooting period (0.756 g/100 g DW), respectively.

Table 1

**Individual catechins
and total phenolic content in seed propagated black tea samples
at different shooting periods, g/100 g DW**

Catechins (g/100 g DW)	Shooting period		
	may	july	september
EGCG	0.941a	0.855b	0.756c
EGC	0.752a	0.640b	0.590c
EC	0.411a	0.367b	0.307b
ECG	0.654a	0.605b	0.510c
CG	0.033 ^{NS}	0.030	0.0206
GCG	0.044a	0.042a	0.032b
C	0.307a	0.278b	0.222c
Total phenolics (g GAE/100 g DW)	7.030a	6.890b	6.450c

Note: There were significant ($P < 0.05$) differences among the different letters in the same lines.

NS: Non Significant

The other individual catechins are also revealed similar patterns (tbl. 1). For example epigallocatechin (EGC) content was also the highest at first shooting period (0.752 g/100 g DW), and it was decreased with prolonged harvest period (0.640 g/100 g DW at second harvest and 0.590 g/100 g DW at last shooting period), respectively.

EC and ECG content were the highest at first shooting period (0.411 and 0.654 g/100 g DW) and decreased to 0.367 and 0.605 g/100 g DW at second shooting period. At third shooting period, those values decreased again to 0.307 and 0.510 g/100 g DW), respectively (tbl. 1).

Ozdemir et al. (2018) reported EGCG, EGC, EC and ECG amounts in black tea samples between 0.819–0.976 g /100 g DW, 0.661–0.785 g/100 g DW, 0.274–0.329 g/100 g DW and 0.437–0.523 g/100 g DW, respectively at three shooting periods. Our results are good agreement with above study. The results also in accordance with previously reported studies by Khokhar and Magnusdottir, (2002) and Serpen et al., (2012).

CG, GCG and C content were found to be the highest at first shooting period (0.033, 0.044 and 0.307 g/100 g DW) while it was the lowest at the last shooting period (0.026, 0.032 and 0.222 g/100 g DW) (tbl. 1).

Catechins and their derivatives are thought to contribute to the beneficial effects ascribed to tea. Tea catechins and polyphenols are effective scavengers

of reactive oxygen species *in vitro* and may also function indirectly as antioxidants through their effects on transcription factors and enzyme activities. The fact that catechins are rapidly and extensively metabolized emphasizes the importance of demonstrating their antioxidant activity *in vivo* [5]. EGCG was the richest catechins. It has been reported that EGCG possessing the greatest number of phenolic hydroxyl groups manifests the strongest antioxidant activity in catechins, which is even stronger than vitamins C and E [14]. It has been reported that EGCG could exhibit multiple bioactivities, especially anticancer effects, through inhibiting cancer stem cells and modulating molecular events associated with cancer cell proliferation, apoptosis, immunity and so on [4, 17].

Total phenolic content were found between 6.45–7.03 g GAE/100 g DW among shooting periods. Ozdemir et al. (2018) found that total phenolic contents of different Turkish black tea samples were in the range of 4.75 to 6.97 g GAE per 100 g. Karadeniz and Koca (2009) reported that Turkish black tea samples had total phenolic content between 7.29–8.39 g GAE per 100 g DW. The total phenolic content of this study were found to be similar to Ozdemir et al. [11] and was lower than those reported previously in the literature.

It is well known that plucking standards and processing conditions could be affect total phenolic content of tea samples. Total phenolic content decreased during the shooting periods, which can be directly related to the total phenolic content of the fresh tea leaves [16].

References

1. Asami D.K., Hong Y.J., Barrett D.M., Mitchell A.E. Comparison of the total phenolic and ascorbic acid content of freeze-dried and air-dried marionberry, strawberry, and corn grown using conventional, organic, and sustainable agricultural practices // *J. Agr Food Chem.* – 2003. – Vol. 51. – P.1237-1241. – doi: 10.1021/jf020635c .
2. Cakmakci R., Erturk Y., Donmez M.F., Erat M., Haznedar A., Sekban R. Tea growth and yield in relation to mixed cultures of N₂ fixing and phosphate solubilizing bacteria // *The Journal of Ege University Faculty of Agriculture.* – Special Issue. – 2012. – Vol. 1. – P. 17-21. – ISSN 1018-8851.
3. FAO. Statistical database: Food and Agriculture Organization of the United Nations, 2019.
4. Gan R.Y., Li H.B., Sui Z.Q., Corke H. Absorption, metabolism, anti-cancer effect and molecular targets of epigallocatechin gallate (EGCG): An updated review // *Crit. Rev. Food Sci.* – 2018. – Vol. 58. – P. 924-941. – doi: 10.1080/10408398.2016.1231168.
5. Higdon J.V., Frei B. Tea catechins and polyphenols: health effects, metabolism, and antioxidant functions // *Crit Rev. Food Sci Nutr.* – 2003. – Vol. 43(1). – P. 89-143. – doi: 10.1080/10408690390826464.
6. Kafkas K., Ercisli S., Doğan Y., Ertürk Y., Haznedar A., Sekban R. Polymorphism and genetic relationships among tea genotypes from turkey revealed by amplified fragment length polymorphism markers // *Journal of American Society Horticultural Sciences.* – 2009. – Vol. 134(4). – P. 428-434. – doi: 10.21273/JASHS.134.4.428.
7. Karadeniz B., Koca I. Phenolic profiles and antioxidant properties of Turkish black tea manufactured with orthodox method // *Asian J Chem.* – 2009. – Vol. 21. – P. 6803-6810. – ISSN 0970-7077.

8. Kelebek H. LC-DAD–ESI-MS/MS characterization of phenolic constituents in Turkish black tea: effect of infusion time and temperature // *Food Chem.* – 2016. – Vol. 204. – P. 227-238. – doi: 10.1016/j.foodchem.2016.02.132.
9. Khokhar S., Magnusdottir S.G.M. Total phenol, catechin, and caffeine contents of teas commonly consumed in the United Kingdom // *J Agr Food Chem.* – 2002. – Vol. 50. – P. 565-570. – doi: 10.1021/jf010153l.
10. Oksuz M. Morphological, yield and quality properties of tea clones in Turkey. – Caykur, Rize, Turkey. – 1987.
11. Ozdemir F., Nadeem H.S., Akdogan A., Dincer C., Topuz A. Effect of altitude, shooting period, and tea grade on the catechins, caffeine, theaflavin, and thearubigin of Turkish black tea // *Turk J Agric For.* – 2018. – Vol. 42. – P. 334-340. – doi: 10.3906/tar-1710-21.
12. Serpen A., Pelvan E., Alasalvar C., Mogol B.A., Yavuz H.T., Gokmen V., Ozcan N., Ozcelik B. Nutritional and functional characteristics of seven grades of black tea produced in Turkey // *J. Agr Food Chem.* – 2012. – Vol. 60. – P. 7682-7689. – doi: 10.1021/jf302058d.
13. Seyis F., Yurteri E., Ozcan A., Savsatli Y. Organic tea production and tea breeding in Turkey: Challenges and Possibilities // *Ekin Journal of Crop Breeding and Genetics.* – 2018. – Vol. 4(1). – P. 60-69. – ISSN 2149-1275.
14. Siddiqui I.A., Afaq F., Adhami V.M., Ahmad N., Mukhtar H. Antioxidants of the beverage tea in promotion of human health // *Antioxid. Redox Signal.* – 2004. – Vol. 6. – P. 571-582. – doi: 10.1089/152308604773934323.
15. Yurteri E., Ozcan A., Seyis F. Tea (*Camellia sinensis*) cultivation and breeding in turkey: Past and Present Status // *Ekin Journal of Crop Breeding and Genetics.* – 2019. – Vol. 5(2). – P. 111-119. – ISSN 2149-1275.
16. Zhang C., Suen C.L.C., Yang C., Quek S.Y. Antioxidant capacity and major polyphenol composition of teas as affected by geographical location, plantation elevation and leaf grade // *Food Chem.* – 2018. – Vol. 244. – P. 109-119. – doi: 10.1016/j.foodchem.2017.09.126.
17. Zhao C.N., Tang G.Y., Cao S.Y., Xu X.Y., Gan R.Y., Liu Q., Mao Q.Q., Shang A., Li H.B. Phenolic profiles and antioxidant activities of 30 tea infusions from green, black, oolong, white, yellow and dark teas // *Antioxidants (Basel).* – 2019. – Vol. 8(7). – P. 215. – doi: 10.3390/antiox8070215.
18. Zuo Y.G., Chen H., Deng Y.W. Simultaneous determination of catechins, caffeine and gallic acids in green, oolong, black and pu-erh teas using HPLC with a photodiode array detector // *Talanta.* – 2002. – Vol. 57. – P. 307-316. – doi: 10.1016/s0039-9140(02)00030-9.

**ИЗМЕНЕНИЕ СОДЕРЖАНИЯ
БИОАКТИВНЫХ ВЕЩЕСТВ В РАЗМНОЖЕННЫХ СЕМЕНАМИ
РАСТЕНИЯХ ЧАЯ В РАЗНЫЕ ПЕРИОДЫ
ПОБЕГООБРАЗОВАНИЯ**

Каратас Н., Эрчисли С., Самарина Л. С.

¹ *Университет им. Ататюрка,
факультет медицинских наук, кафедра питания и диетологии,
г. Эрзурум, Турция*

² *Университет им. Ататюрка,
Факультет растениеводства, кафедра садоводства,
г. Эрзурум, Турция, e-mail: sercisli@gmail.com*

³ Федеральное государственное бюджетное учреждение науки
«Федеральный исследовательский центр «Субтропический научный центр
Российской академии наук», Отдел биотехнологии,
г. Сочи, Россия, e-mail: q1111w2006@ya.ru

Производство чая крайне важно для фермеров Черноморского региона, поскольку данная культура является доминирующим сельскохозяйственным продуктом. Основная часть чайных садов региона была размножена фермерами, использующими семенной метод на протяжении последних 50 лет, что привело к большим вариациям большинства характеристик растений. В настоящем исследовании описано влияние различных периодов побегообразования на содержание биологически активных веществ чайного растения, размноженного семенами. Результаты показали значительные различия между большинством искомых параметров по периодам побегообразования. Общее содержание фенолов было самым высоким в первый период побегообразования (6,45–7,03 г эквивалента галловой кислоты/100 г) по сухой массе. Среди катехинов в образцах чёрного чая доминировал эпигаллокатехин галлат (EGCG) (0,756–0,941 г/100 г сырой массы), за которым следовали эпигаллокатехин (0,590–0,752 г/100 г сырой массы) и эпикатехин галлат (0,510–0,654 г/100 г сырой массы), соответственно. В целом в первый период побегообразования (май) содержание катехинов было наибольшим, что улучшило качество чая, а с последующими периодами побегообразования содержание катехинов неуклонно снижалось. Общее содержание фенола было в пределах 6,45–7,03 г ГК/100 г ДВ в разные периоды побегообразования. Результаты показали, что в первый период побегообразования образцы чёрного чая имели более высокое содержание биологически активных веществ.

Ключевые слова: чай, разнообразие, фитохимические вещества, семенное размножение.

УДК 633.72:581.19

doi: 10.31360/2225-3068-2020-75-74-81

ОБЩЕЕ СОДЕРЖАНИЕ АНТОЦИАНОВ В 3-ЛИСТНЫХ ФЛЕШАХ И ИЗМЕНЕНИЕ ИХ КОЛИЧЕСТВА ПРИ ПЕРЕРАБОТКЕ

Платонова Н. Б., Белоус О. Г.

*Федеральное государственное бюджетное учреждение науки
«Федеральный исследовательский центр «Субтропический научный центр
Российской академии наук»,
г. Сочи, Россия, e-mail: oksana191962@mail.ru*

Проведены исследования по определению антоцианов в 3-листной флешке, зелёном и чёрном чае. Определены основные закономерности в их накоплении по месяцам. Наибольшее количество антоцианов накапливается в июле (308,5 г/100 г), наименьшее – в мае (150,2 г/100 г), что связано с длительным засушливым периодом, сопровождаемым высокими температурами воздуха.